

CPH-A213.1292

HUPD-9216

December 1992

Conformal dynamics of quantum gravity with torsion

I. ANTONIADIS

Centre de Physique Théorique

Ecole Polytechnique

91128 Palaiseau, France

and

S.D. ODINTSOV[★]

Department of Physics

Hiroshima University

Higashi-Hiroshima 724, Japan

ABSTRACT

The trace anomaly induced dynamics of the conformal factor is investigated in four-dimensional quantum gravity with torsion. The constraints for the coupling constants of torsion matter interaction are obtained in the infrared stable fixed point of the effective scalar theory.

★ On leave from Pedagogical Institute, 634041 Tomsk, Russia

In a recent paper [1] the trace anomaly induced dynamics of the conformal factor in four-dimensional quantum gravity has been investigated. The conformal sector was described by spacetime metrics of the form:

$$g_{\mu\nu}(x) = e^{2\sigma(x)}\eta_{\mu\nu} \ ,$$

where e^σ is the conformal factor and $\eta_{\mu\nu}$ is the flat metric. The generalization to curved fiducial metric was considered in refs.[2,3]. The effective action for σ can be obtained by integrating the general form of trace anomaly in curved spacetime [4,3]. It turns out that the resulting action is local in σ and possesses a non-trivial, infrared stable critical point which can be computed to all orders in perturbation theory [1]. In particular, the anomalous scaling dimension of the conformal field e^σ was found using the same line of reasoning as in two-dimensional gravity [5]. It was argued that the effective σ theory describes four-dimensional quantum gravity at large distances and provides a framework for a dynamical solution of the cosmological constant problem [1].

The purpose of this note is to generalize the results of ref.[1] in the case of quantum gravity with torsion. Torsion interactions start becoming non-trivial in four dimensions. In fact in two dimensions there is no minimal coupling of matter with torsion. Furthermore, in the non-minimal case [6] torsion appears quadratically and, after functional integration, 2d induced gravity with torsion is equivalent to 2d gravity without torsion up to coupling constants redefinitions. In four dimensions minimal interactions of spinors with torsion are non-trivial. Moreover, the torsion dependence of trace anomaly is much more complicated than in two dimensions [7], so that one may expect to find some new properties of the conformal factor dynamics in this case.

Let us start with the free actions for scalar and spinor fields in curved spacetime in the presence of torsion $T^\alpha_{\gamma\beta}$. Recall that in a theory with torsion the connection $\Gamma^\alpha_{\beta\gamma}$ is not symmetric in the lower indices:

$$\Gamma^\alpha_{\beta\gamma} - \Gamma^\alpha_{\gamma\beta} = T^\alpha_{\gamma\beta} \neq 0. \tag{1}$$

This relation is the origin of torsion. Here, we consider the case where the only non-vanishing components of torsion are those which correspond to the totally antisymmetric part, $S^\mu \equiv i\varepsilon^{\alpha\beta\nu\mu}T_{\alpha\beta\nu}$, called pseudotrace. This is the only part of torsion which can interact minimally with matter (see ref.[7] for more details).

The conformally invariant quadratic actions for scalars (S_0) and spinors ($S_{\frac{1}{2}}$) are [7]:

$$S_0 = \frac{1}{2} \int d^4x \sqrt{-g} \{ g^{\alpha\beta} \partial_\alpha \varphi \partial_\beta \varphi + \frac{1}{6} R \varphi^2 + \xi_0 S_\mu S^\mu \varphi^2 \}, \quad (2)$$

$$S_{\frac{1}{2}} = i \int d^4x \sqrt{-g} \{ \bar{\psi} (\gamma^\mu \nabla_\mu - \xi_{\frac{1}{2}} \gamma_5 \gamma^\mu S_\mu) \psi \}, \quad (3)$$

where $\xi_0, \xi_{\frac{1}{2}}$ are arbitrary constants. These actions are invariant under the conformal transformations:

$$g_{\mu\nu} \rightarrow e^{2\sigma} g_{\mu\nu}, \quad \varphi \rightarrow e^{-\sigma} \varphi, \quad \psi \rightarrow e^{-\frac{3\sigma}{2}} \psi \quad (4)$$

for any values of $\xi_0, \xi_{\frac{1}{2}}$. The choice $\xi_0 = 0, \xi_{\frac{1}{2}} = \frac{1}{8}$ corresponds to the minimal interaction of matter with torsion. Here we do not consider torsion interaction with gauge fields. Such interaction is non minimal, and it exists in general only in the abelian case leading to terms which are first order in quantum fields [7]. It is therefore irrelevant for our purposes.

The general form of the trace anomaly for free conformally invariant matter in curved spacetime with torsion is:

$$\begin{aligned} T^\mu_\mu = & b(F + \frac{2}{3} \square R) + b' G + b'' \square R + a_1 F_{\mu\nu} F^{\mu\nu} \\ & + a_2 (S_\mu S^\mu)^2 + a_3 \square (S_\mu S^\mu) + a_4 \nabla_\mu (S_\nu \nabla^\nu S^\mu - S^\mu \nabla_\nu S^\nu), \end{aligned} \quad (5)$$

where

$$F \equiv C_{\mu\nu\alpha\beta} C^{\mu\nu\alpha\beta}, \quad G = R^2_{\mu\nu\alpha\beta} - 4R^2_{\mu\nu} + R^2, \quad F_{\mu\nu} = \nabla_\mu S_\nu - \nabla_\nu S_\mu,$$

$$\begin{aligned}
b &= \frac{1}{120(4\pi)^2}(N_0 + 6N_{\frac{1}{2}} + 12N_1), \quad b' = -\frac{1}{360(4\pi)^2}(N_0 + 11N_{\frac{1}{2}} + 62N_1), \\
a_1 &= -\frac{2}{3(4\pi)^2} \sum \xi_{\frac{1}{2}}^2, \quad a_2 = \frac{1}{2(4\pi)^2} \sum \xi_0^2, \quad a_3 = \frac{1}{3(4\pi)^2} \sum (2\xi_{\frac{1}{2}}^2 - \frac{1}{2}\xi_0^2), \\
a_4 &= -\frac{2}{3(4\pi)^2} \sum \xi_{\frac{1}{2}}^2,
\end{aligned}$$

with N_0 , $N_{\frac{1}{2}}$ and N_1 the number of scalars, Dirac fermions and vectors. The coefficients b and b' can be found in ref.[8], b'' is arbitrary since it can be changed by adding a local R^2 term in the action, while the coefficients a_1 , a_2 , a_3 and a_4 are given in ref.[7] for the case of minimal interaction with torsion ($\xi_0 = 0$, $\xi_{\frac{1}{2}} = \frac{1}{8}$) $a_2 = 0$.

We now choose the conformal parametrization $g_{\mu\nu} = e^{2\sigma(x)}\bar{g}_{\mu\nu}$ and $S_\mu = \bar{S}_\mu$, where $\bar{g}_{\mu\nu}$ and $\bar{S}_\mu$ are the fiducial metric and torsion, respectively. The conformal factor acquires dynamics by an effective action whose σ -variation reproduces the anomalous trace (5):

$$\begin{aligned}
T_\mu^\mu &= bF + b'(G - \frac{2}{3}\square R) + [b'' + \frac{2}{3}(b + b'')] \square R + a_1 F_{\mu\nu} F^{\mu\nu} \\
&\quad + a_2 (S_\mu S^\mu)^2 + a_3 \square (S_\mu S^\mu) + a_4 \nabla_\mu (S_\nu \nabla^\nu S^\mu - S^\mu \nabla_\nu S^\nu) \\
&\equiv \frac{1}{\sqrt{-g}} \frac{\delta}{\delta \sigma(x)} S_{anom}(\sigma)
\end{aligned} \tag{6}$$

Integrating eq. (6) we get (see refs.[1,5] for $S_\mu = 0$, and ref.[7] for $S_\mu \neq 0$) :

$$\begin{aligned}
S_{anom}[\sigma] &= \int d^4x \sqrt{-\bar{g}} \{ \sigma [b\bar{F} + b'(\bar{G} - \frac{2}{3}\bar{\square}\bar{R}) + a_1\bar{F}_{\mu\nu}^2 + a_2\bar{S}^4] \\
&\quad + 2b'\sigma\bar{\Delta}\sigma - a_3\bar{\nabla}_\mu\sigma\bar{\nabla}^\mu\bar{S}^2 - a_4\bar{\nabla}_\mu\sigma(\bar{S}_\nu\bar{\nabla}^\nu\bar{S}^\mu - \bar{S}^\mu\bar{\nabla}_\nu\bar{S}^\nu) \\
&\quad + (a_3 + \frac{1}{2}a_4)\bar{S}^2\bar{\nabla}_\mu\sigma\nabla^\mu\sigma + a_4\bar{S}^\mu\bar{S}^\nu\bar{\nabla}_\mu\sigma\bar{\nabla}_\nu\sigma \\
&\quad - \frac{1}{12}[b'' + \frac{2}{3}(b + b'')] \int d^4x \sqrt{-\bar{g}} [\bar{R} - 6\bar{\square}\sigma - 6(\bar{\nabla}_\mu\sigma)(\bar{\nabla}^\mu\sigma)]^2 \},
\end{aligned} \tag{7}$$

where $\bar{S}^2 = \bar{S}_\mu\bar{S}^\mu$, and Δ is the conformally covariant fourth-order operator acting on scalars,

$$\Delta = \square^2 + 2R^{\mu\nu}\nabla_\mu\nabla_\nu - \frac{2}{3}R\square + \frac{1}{3}\nabla_\mu R\nabla^\mu. \tag{8}$$

In eq. (7) we have omitted σ -independent terms.

Now as in two dimensions [5] we add S_{anom} to the classical gravity action:

$$\begin{aligned} S_{cl} &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} (R + h S_\mu S^\mu - 2\Lambda) \\ &= \frac{1}{2\kappa} \int d^4x \sqrt{-\bar{g}} \{ e^{2\sigma} [R + 6(\bar{\nabla}_\mu \sigma)(\bar{\nabla}^\mu \sigma)] - 2\Lambda e^{4\sigma} + h e^{2\sigma} \bar{S}^2 \}, \end{aligned} \quad (9)$$

where h is a constant parameter (e.g. for the Einstein-Cartan theory $h = -\frac{1}{24}$). Thus, the total effective action is:

$$S_{eff} = S_{anom} + S_{cl} \quad (10)$$

When the conformal factor is quantized, the requirement of general covariance in the vacuum state of σ -theory implies the vanishing of the total trace at the quantum level [3]. Therefore, the β -functions of all couplings appearing in the effective σ -action must vanish. Without loss of generality, let us consider the effective theory (10) with flat fiducial metric and arbitrary constant torsion background S_μ :

$$\begin{aligned} S_{eff} &= -\frac{Q^2}{(4\pi)^2} \int d^4x (\Box \sigma)^2 - \zeta \int d^4x [2\alpha (\partial_\mu \sigma)^2 \Box \sigma + \alpha^2 (\partial_\mu \sigma)^4] \\ &\quad + \gamma \int d^4x e^{2\alpha\sigma} (\partial_\mu \sigma)^2 - \frac{\lambda}{\alpha^2} \int d^4x e^{4\alpha\sigma} + \int d^4x \{ (a_3 + \frac{1}{2}a_4) S^2 (\partial_\mu \sigma)^2 \\ &\quad + a_4 S^\mu S^\nu \partial_\mu \sigma \partial_\nu \sigma + \frac{h}{2\kappa\alpha^2} e^{2\alpha\sigma} S^2 + a_2 \sigma S^4 \}, \end{aligned} \quad (11)$$

where the transformations $\sigma \rightarrow \sigma\alpha$, $S_{eff} \rightarrow \alpha^{-2} S_{eff}$ are made, and the notations of ref.[1] are introduced,

$$\frac{Q^2}{(4\pi)^2} = 2b + 3b', \quad \zeta = 2b + 2b' + 3b'', \quad \gamma = \frac{3}{\kappa}, \quad \lambda = \frac{\Lambda}{\kappa}. \quad (12)$$

α is the anomalous scaling dimension of the conformal factor at the non-trivial infrared critical point we want to determine.

A simple power counting argument shows that the effective action (11) is renormalizable in σ -perturbation theory if we add the action of the external torsion-field:

$$S_{ext} = \frac{\eta}{\alpha^2} \int d^4x (S_\mu S^\mu)^2. \quad (13)$$

The ζ coupling renormalization does not depend on the remaining interactions of (11), and a straight-forward one-loop calculation of its β -function shows that $\zeta = 0$ is an infrared stable fixed point [1]. At $\zeta = 0$, the contribution to the anomaly coefficients b and b' of the σ -field itself is coming from the quartic operator Δ (8) and was computed in ref.[3]. Moreover, due to the quartic kinetic operator, the effective action (11) becomes superrenormalizable and all β -functions can be computed exactly. The presence of torsion background does not modify the β -functions of γ and λ couplings. Their vanishing imply [1]:

$$\alpha_{\pm} = \frac{1 \pm \sqrt{1 - 4/Q^2}}{2/Q^2}, \quad \frac{\lambda}{\gamma^2} = \frac{2\pi^2}{Q^2} \left[1 + \frac{4\alpha^2}{Q^2} + \frac{6\alpha^4}{Q^4} \right]. \quad (15)$$

Eq. (15) determines at the fixed point the anomalous scaling dimension α in terms of the “central charge” Q^2 , and relates the cosmological with the Newtonian couplings. Now we can calculate the additional β -functions connected with the torsion.

We first perform a one-loop computation using the standard algorithm [9] and then we will show that the one-loop result is exact. Expanding σ around a classical background solution and keeping only terms which are quadratic in quantum fluctuations, we can write the resulting kinetic operator corresponding to the action (11) in the following form:

$$\hat{H} = \square^2 + D^{\mu\nu} \nabla_\mu \nabla_\nu + H^\mu \nabla_\mu + p \quad (16)$$

where H^μ does not give any contribution to the one-loop divergences and can be

dropped,

$$D^{\mu\nu} = \frac{(4\pi)^2}{Q^2} [\gamma e^{2\alpha\sigma} g^{\mu\nu} + (a_3 + \frac{1}{2}a_4) S^2 g^{\mu\nu} + a_4 S^\mu S^\nu],$$

$$p = -\frac{(4\pi)^2}{Q^2 \kappa} h e^{2\alpha\sigma} S^2 - 2(\frac{4\pi}{Q})^2 \gamma \alpha^2 e^{2\alpha\sigma} (\partial_\mu \sigma)^2 + 8\lambda (\frac{4\pi}{Q})^2 e^{4\alpha\sigma}.$$

The corresponding a_2 Schwinger-De Witt coefficient in the heat kernel expansion of $\hat{H}$ is:

$$\begin{aligned} a_2(\hat{H}) &= \frac{1}{(4\pi)^2} \left\{ -p + \frac{1}{48} D^2 + \frac{1}{24} D_{\mu\nu} D^{\mu\nu} \right\} \\ &= e^{2\alpha\sigma} S^2 \left[\frac{h}{\kappa Q^2} + \frac{\gamma(4\pi)^2}{Q^4} (a_3 + \frac{3}{4}a_4) \right] + S^4 \frac{(4\pi)^2}{Q^4} \left[\frac{1}{2}a_3^2 + \frac{3}{4}a_3 a_4 + \frac{5}{16}a_4^2 \right], \end{aligned} \quad (17)$$

where we neglected total derivatives, like $\square D$ and $\partial_\mu \partial_\nu D^{\mu\nu}$, as well as S independent terms which have been already discussed. Taking also into account the “classical” contribution due to the change of the scaling dimension of the conformal factor e^σ we get:

$$\begin{aligned} \beta_h &= (2 - 2\alpha + \frac{2\alpha^2}{Q^2}) \frac{h}{2\kappa} + \frac{\alpha^2 \gamma (4\pi)^2}{Q^4} (a_3 + \frac{3}{4}a_4), \\ \beta_\eta &= \frac{\alpha^2 (4\pi)^2}{Q^4} \left[\frac{1}{2}a_3^2 + \frac{3}{4}a_3 a_4 + \frac{5}{16}a_4^2 \right]. \end{aligned} \quad (18)$$

One can now show by a Feynman graph analysis similar to the one done in ref.[1] that β_h is the exact β -function and there are no higher order corrections. In fact, all additional terms in (11) involving the torsion background amount only to a change of σ -propagator, with the exception of the h -term which generates new vertices. A simple power counting argument then shows that the only graphs with primitive divergences must contain at most two γ -vertices or exactly one λ or h -vertex. It follows that the S -dependent part of divergences come from tadpole diagrams with one h or γ -vertex and lead to the expression for β_h in (18). β_η is associated to vacuum (σ -independent) terms which can be made zero by an appropriate σ -translation, due to the presence of the last linear term in σ in (11).

The gravitational (spin-two and reparametrization ghost) contribution remains an open question. In ref.[3], it was argued that in the infrared this amounts only to a change of the trace-anomaly coefficients (5) in the effective σ -theory. Furthermore, this contribution was calculated at the one-loop level using the Einstein action, as well as the conformally invariant but four-derivative Weyl tensor squared action. In the presence of non-vanishing torsion the computation becomes more complicated. A potentially interesting case could be the example of higher derivative gravity with torsion which seems to avoid the usual unitarity problem for some values of the parameters at the classical level [10].

Focussing in the σ -field sector, and using eq. (15), the vanishing of β_h gives

$$a_3 + \frac{3}{4}a_4 = 0. \tag{19}$$

This condition constraints the torsion parameters of the theory at the infrared stable fixed point. In the case of pure matter it implies $\xi_0 = \xi_{\frac{1}{2}}$, which for minimal interaction is satisfied only for zero torsion.

Acknowledgements: This work was supported in part by the EEC contracts SC1-0394-C and SC1-915053. SDO wishes to thank Particle Theory Group at Hiroshima University for kind hospitality and JSPS for financial support. We also thank I.L. Shapiro for discussions on the point that β_h is given by the one-loop result.

REFERENCES

1. I. Antoniadis and E. Mottola, Phys. Rev. **D45** (1992) 2013.
2. S.D. Odintsov. Z. Phys. **C54** (1992), 531.
3. I. Antoniadis, P.O. Mazur and E. Mottola, Conformal Symmetry and central charges in 4 dimensions, LA-UR 92 preprint.
4. R.J. Reigert, Phys. Lett. **B134** (1984) 56; E.T. Tomboulis, Nucl. Phys. **B329** (1990) 410; S.D. Odintsov and I.L. Shapiro, Class. Quant. Grav. **8** (1991) L57.
5. A.M. Polyakov, Mod. Phys. Lett. **A2** (1987) 899; V.G. Knizhnik, A.M. Polyakov and A.B. Zamolodchikov, Mod. Phys. Lett. **A3** (1988) 819; J. Distler and H. Kawai, Nucl. Phys. **B321** (1989) 509; F. David, Mod. Phys. Lett. **A3** (1988) 1651.
6. H. Leutwyler, Phys. Lett. **153** (1985) 65; L. Alvarez - Gaumé and P. Ginsparg, Ann. Phys. (NY) **161** (1985) 423; K. Li, Phys. Rev. **D34** (1986) 2292; T. Fukuyama and K. Kamimura, Phys. Lett. **B200** (1988) 75; S.D. Odintsov, Rivista Nuovo Cim. **15** (1992) No.3 .
7. I.L. Buchbinder, S.D. Odintsov and I.L. Shapiro, Phys. Lett. **162** (1985) 93; Rivista Nuovo Cim. **12** (1989) 1; see also Effective action in quantum gravity, IOP Publishing, 1992 (Chapter 4,5).
8. N. Birrell and P. Davies, Quantum fields in curved space, CUP, 1982 and references therein.
9. E.S. Fradkin and A.A. Tseytlin, Nucl. Phys. **B201** (1982) 469.
10. E. Sezgin and P. van Nieuwenhuizen, Phys. Rev. **D21** (1980) 3269.